

University of Groningen

Analysis of bacterial adhesion forces

Chen, Yun

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chen, Y. (2014). *Analysis of bacterial adhesion forces: a study using atomic force microscopy*. [Thesis fully internal (DIV), University of Groningen]. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 1

General Introduction

Bacteria adhere to virtually all natural and man-made surfaces. Initiated by their adhesion, micro-organisms grow on a surface and form complex multi-cellular structures, referred to as biofilms.^{1,2} In biofilms, micro-organisms are protected against environmental attacks,³⁻⁵ such as antimicrobials, and therefore biofilms are often the cause of persistent contamination or infection.⁶⁻⁸ Studies on screening of different preventive measures and removal of biofilms⁹ are gaining significant attention in various fields of application, ranging from food and water processing¹⁰ to modern medicine¹¹ and dentistry (Fig. 1).¹²




Figure 1. Common sites of primary and secondary (highlighted in green) biofilm infection in the human body. Source: Center for Biofilm Engineering, Montana State University-Bozeman.

Surface modification can be effective in reducing microbial adhesion to substratum surfaces,¹³ while, interestingly, the susceptibility of a biofilm for antimicrobials in solution varies depending on the properties of the substratum surface. Different classes of antimicrobials, such as quaternary ammonium compounds or antimicrobial peptides, remain antimicrobially active when immobilized to a substratum surface.^{14,15} Hypothetically, different generic mechanisms of action must exist for antimicrobials in solution and immobilized on a surface.

Insight into mechanisms of biofilm formation can lead to improved preventive mechanisms and requires knowledge of initial bacterial adhesion and the surface properties of the adhering bacteria.

BACTERIAL RESPONSE TO THEIR ADHESION ON A SUBSTRATUM SURFACE

It has been recently hypothesized¹⁶ that an adhering bacterium responds to a substratum surface depending on the strength of the force by which it adheres. In this respect, three regimes of adhesion forces have been proposed (Fig. 2):

1. A “planktonic” regime, where the adhesion force is very weak so that the bacteria hardly sense that they are on a surface and therefore remain in their planktonic state,^{17–19} in which they are susceptible to antimicrobials;
2. An “interaction” regime, where adhesion forces induce a cascade of genotypic and phenotypic changes²⁰ that render the organism more resistant to antimicrobial agents;^{21,22}
3. A “lethal regime” in which strong adhesion forces de-activate adhering bacteria to impede growth and cause cell death.^{16,23}




Figure 2. Three regimes of bacterial adhesion forces to substratum surfaces that dictate the bacterial response to a surface. (Illustration adapted from Busscher and Van der Mei¹⁶)

In the different regimes of adhesion forces, it can be envisaged that a bacterium will undergo different degrees of cell wall deformation to which the organism subsequently responds. The term “stress-deactivation” has been coined for this hypothetical phenomenon.²³ Accordingly, knowledge of the forces by which bacteria adhere to a surface are of pivotal importance in understanding further events during biofilm formation.

BACTERIAL ADHESION FORCES USING ATOMIC FORCE MICROSCOPY (AFM)

Bacterial adhesion to surfaces can be assessed using various biochemical and physico-chemical approaches.^{24,25} Among these methods, AFM, right after its invention in the 1970s,^{26,27} showed most promising.^{28,29} The distance dependence

of the interaction force between a bacterium and a substratum surface can be recorded using AFM as a force-distance curve with an approach and retract stage (Fig. 3). The magnitude of the adhesion force is usually determined by the rupture force when detaching the AFM cantilever, equipped with a bacterial probe, from the substratum surface.




Figure 3. Illustration of measuring bacterial interaction forces using AFM and an example of the approach and retract force-distance curves.

Theoretical models based on the Derjaguin-Landau-Verwey-Overbeek (DLVO) theory^{30,31} and its extended version (the X-DLVO theory)³² are widely applied to derive the long-range (*e.g.* Lifshitz-Van der Waals and electrical double-layer interactions) and short-range (*e.g.* Lewis acid-base interactions) contributions to the total adhesion force, based on measured zeta potentials and contact angles

using various liquids.³² The statistics-based Poisson analysis of bacterial adhesion forces measured using AFM provides a more experimental alternative to distinguish between long-range and short-range contributions to the overall interaction force.^{33,34} However, these methods do not always concur in the magnitude and even the attractive or repulsive nature of the short-range interaction.^{35,36} It has been argued that (X)DLVO-based models fail to take microscopic bacterial cell wall structures into consideration, whereas Poisson analysis of AFM-based adhesion forces includes effects of the microscopic features of the interface between a bacterium and a substratum surface.

AIM OF THIS STUDY

The aim of this thesis is to determine bacterial adhesion forces to substratum surfaces and associated cell wall deformation and evaluate their role in bacterial susceptibility to antimicrobials, either in solution or immobilized to a substratum surface.

REFERENCES

1. Fletcher, M. Bacterial biofilms and biofouling. *Curr. Opin. Biotechnol.* **5**, 302–306 (1994).
2. Berk, V., Fong, J. C. N., Dempsey, G. T., Develioglu, O. N., Zhuang, X., Liphardt, J., Yildiz, F. H. & Chu, S. Molecular architecture and assembly principles of *Vibrio cholerae* biofilms. *Science* **337**, 236–239 (2012).
3. Stewart, P. S. & Costerton, J. W. Antibiotic resistance of bacteria in biofilms. *Lancet* **358**, 135–138 (2001).
4. Mah, T.-F., Pitts, B., Pellock, B., Walker, G. C., Stewart, P. S. & O'Toole, G. A. A genetic basis for *Pseudomonas aeruginosa* biofilm antibiotic resistance. *Nature* **426**, 306–310 (2003).
5. Anderson, G. G. & O'Toole, G. A. Innate and induced resistance mechanisms of bacterial biofilms. *Curr. Top. Microbiol. Immunol.* **322**, 85–105 (2008).
6. Costerton, J. W., Stewart, P. S. & Greenberg, E. P. Bacterial biofilms: a common cause of persistent infections. *Science* **284**, 1318–1322 (1999).
7. Parsek, M. R. & Singh, P. K. Bacterial biofilms: an emerging link to disease pathogenesis. *Annu. Rev. Microbiol.* **57**, 677–701 (2003).
8. Hall-Stoodley, L., Costerton, J. W. & Stoodley, P. Bacterial biofilms: from the natural environment to infectious diseases. *Nat. Rev. Microbiol.* **2**, 95–108 (2004).
9. Vejborg, R. M. & Klemm, P. Blocking of bacterial biofilm formation by a fish protein coating. *Appl. Environ. Microbiol.* **74**, 3551–3558 (2008).
10. Hu, X., Xu, K., Wang, Z., Ding, L. & Ren, H. Characteristics of biofilm attaching to carriers in moving bed biofilm reactor used to treat vitamin C wastewater. *Scanning* **35**, 283–291 (2013).
11. Schaer, T. P., Stewart, S., Hsu, B. B. & Klibanov, A. M. Hydrophobic polycationic coatings that inhibit biofilms and support bone healing during infection. *Biomaterials* **33**, 1245–1254 (2012).
12. Traverse, C. C., Mayo-Smith, L. M., Poltak, S. R. & Cooper, V. S. Tangled bank of experimentally evolved *Burkholderia* biofilms reflects selection during chronic infections. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E250–E259 (2013).

13. Cheng, G., Zhang, Z., Chen, S., Bryers, J. D. & Jiang, S. Inhibition of bacterial adhesion and biofilm formation on zwitterionic surfaces. *Biomaterials* **28**, 4192–4199 (2007).
14. Fuchs, A. D. & Tiller, J. C. Contact-active antimicrobial coatings derived from aqueous suspensions. *Angew. Chem.-Int. Ed.* **45**, 6759–6762 (2006).
15. Green, J.-B. D., Bickner, S., Carter, P. W., Fulghum, T., Luebke, M., Nordhaus, M. A. & Strathmann, S. Antimicrobial testing for surface-immobilized agents with a surface-separated live–dead staining method. *Biotechnol. Bioeng.* **108**, 231–236 (2011).
16. Busscher, H. J. & Van der Mei, H. C. How do bacteria know they are on a surface and regulate their response to an adhering state? *PLoS Pathog.* **8**, e1002440 (2012).
17. Haldar, J., An, D. Q., de Cienfuegos, L. A., Chen, J. Z. & Klibanov, A. M. Polymeric coatings that inactivate both influenza virus and pathogenic bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 17667–17671 (2006).
18. Ignatova, M., Voccia, S., Gabriel, S., Gilbert, B., Cossement, D., Jérôme, R. & Jérôme, C. Stainless steel grafting of hyperbranched polymer brushes with an antibacterial activity: synthesis, characterization, and properties. *Langmuir* **25**, 891–902 (2009).
19. Hook, A. L., Chang, C.-Y., Yang, J., Luckett, J., Cockayne, A., Atkinson, S., Mei, Y., Bayston, R., Irvine, D. J., Langer, R., Anderson, D. G., Williams, P., Davies, M. C. & Alexander, M. R. Combinatorial discovery of polymers resistant to bacterial attachment. *Nat. Biotechnol.* **30**, 868–875 (2012).
20. Lin, C.-H., Kabrawala, S., Fox, E. P., Nobile, C. J., Johnson, A. D. & Bennett, R. J. Genetic control of conventional and pheromone-stimulated biofilm formation in *Candida albicans*. *PLoS Pathog* **9**, e1003305 (2013).
21. Chiang, W.-C., Nilsson, M., Jensen, P. Ø, Højby, N., Nielsen, T. E., Givskov, M. & Tolker-Nielsen, T. Extracellular DNA shields against aminoglycosides in *Pseudomonas aeruginosa* biofilms. *Antimicrob. Agents Chemother.* **57**, 2352–2361 (2013).
22. Savage, V. J., Chopra, I. & O’Neill, A. J. Population diversification in *Staphylococcus aureus* biofilms may promote dissemination and persistence. *PLoS ONE* **8**, e62513 (2013).

23. Liu, Y., Strauss, J. & Camesano, T. A. Adhesion forces between *Staphylococcus epidermidis* and surfaces bearing self-assembled monolayers in the presence of model proteins. *Biomaterials* **29**, 4374–4382 (2008).
24. Habimana, O., Meyrand, M., Meylheuc, T., Kulakauskas, S. & Briandet, R. Genetic features of resident biofilms determine attachment of *Listeria monocytogenes*. *Appl. Environ. Microbiol.* **75**, 7814–7821 (2009).
25. Rohde, H., Frankenberger, S., Zähringer, U. & Mack, D. Structure, function and contribution of polysaccharide intercellular adhesin (PIA) to *Staphylococcus epidermidis* biofilm formation and pathogenesis of biomaterial-associated infections. *Eur. J. Cell Biol.* **89**, 103–111 (2010).
26. Young, R., Ward, J. & Scire, F. The topografiner: an instrument for measuring surface microtopography. *Rev. Sci. Instrum.* **43**, 999–1011 (1972).
27. Binnig, G., Quate, C. F. & Gerber, C. Atomic force microscope. *Phys. Rev. Lett.* **56**, 930–933 (1986).
28. Allison, D. P., Mortensen, N. P., Sullivan, C. J. & Doktycz, M. J. Atomic force microscopy of biological samples. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* **2**, 618–634 (2010).
29. Lower, S. K. Atomic force microscopy to study intermolecular forces and bonds associated with bacteria. *Adv. Exp. Med. Biol.* **715**, 285–299 (2011).
30. Derjaguin, B. & Landau, L. Theory of the stability of strongly charged lyophobic sols and of the adhesion of strongly charged particles in solutions of electrolytes. *Prog. Surf. Sci.* **43**, 30–59 (1941).
31. Verwey, E. J. W., Overbeek, J. T. G. & Van Nes, K. *Theory of the stability of lyophobic colloids: the interaction of sol particles having an electric double layer*. (Elsevier Publishing Company, 1948).
32. Van Oss, C. J. Acid-base interfacial interactions in aqueous media. *Colloids Surf. A Physicochem. Eng. Asp.* **78**, 1–49 (1993).
33. Han, T., Williams, J. & Beebe, T. Chemical-bonds studied with functionalized atomic-force microscopy tips. *Anal. Chim. Acta* **307**, 365–376 (1995).

34. Williams, J., Han, T. & Beebe, T. Determination of single-bond forces from contact force variances in atomic force microscopy. *Langmuir* **12**, 1291–1295 (1996).
35. Méndez-Vilas, A., Gallardo-Moreno, A. M., Calzado-Montero, R. & González-Martín, M. L. AFM probing in aqueous environment of *Staphylococcus epidermidis* cells naturally immobilised on glass: physico-chemistry behind the successful immobilisation. *Colloids Surf. B Biointerfaces* **63**, 101–109 (2008).
36. Dorobantu, L. S., Bhattacharjee, S., Foght, J. M. & Gray, M. R. Analysis of force interactions between AFM tips and hydrophobic bacteria using DLVO theory. *Langmuir* **25**, 6968–6976 (2009).